


Listing of Claims:

Claim 1 (currently amended): A compound of formula (I):



wherein:

groups R³ and R⁴ are attached to the Z¹ ring structure and groups R⁵ and R⁶ are attached to the Z² ring structure, and n = 1, 2 or 3;

Z¹ and Z² independently represent the carbon atoms necessary to complete a one ring, or two-fused ring aromatic system;


at least one of groups R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ is the group –E–F where E is a single bond or a spacer group having a chain from 1–20 linked atoms selected from the group consisting of carbon, nitrogen and oxygen atoms and F is a target bonding group;

wherein F is either:

- (i) a reactive group selected from carboxyl, succinimidyl ester, sulpho-succinimidyl ester, isothiocyanate, maleimide, haloacetamide, acid halide, hydrazide, vinylsulphone, dichlorotriazine and phosphoramidite; or
- (ii) a functional group selected from hydroxy, amino, sulphydryl, carbonyl (including aldehyde and ketone) and thiophosphate; or

(iii) an affinity tag;

one or more of groups R^{11} , R^{12} , R^{13} and R^{14} are independently selected from the group $-(CH_2)_k-W$, where W is sulphonic acid or phosphonic acid and k is an integer from 1 to 10; when any of groups R^1 and R^2 is not said group $-E-F$, said remaining groups R^1 and R^2 are independently selected from C_1-C_6 alkyl, benzyl either unsubstituted or substituted with sulphonic acid, and the group $-(CH_2)_k-W$, where W and k are hereinbefore defined; when any of groups R^3 , R^4 , R^5 and R^6 is not said group $-E-F$, said remaining groups R^3 , R^4 , R^5 and R^6 are independently selected from hydrogen and sulphonic acid; when any of groups R^{11} , R^{12} , R^{13} and R^{14} is not said group $-(CH_2)_k-W$, said remaining groups R^{11} , R^{12} , R^{13} and R^{14} are independently C_1-C_6 alkyl; remaining groups R^7 are hydrogen or two of R^7 together with the group,



form a hydrocarbon ring system having 5 or 6 atoms.

Claim 2 (original): A compound according to claim 1 wherein at least two of R^{11} , R^{12} , R^{13} and R^{14} are independently $-(CH_2)_k-W$ wherein W and k are hereinbefore defined.

Claim 3 (original): A compound according to claim 1 wherein one of groups R^{11} and R^{12} and one of groups R^{13} and R^{14} is the group $-(CH_2)_k-W$ wherein W and k are hereinbefore defined; and remaining groups R^{11} or R^{12} and R^{13} or R^{14} are C_1-C_6 alkyl.

Claim 4 (previously presented): A compound according to claim 1 wherein W is sulphonic acid.


Claim 5 (previously presented): A compound according to claim 1 wherein $-(CH_2)_k-W$ is selected from $-(CH_2)_3-SO_3H$ and $-(CH_2)_4-SO_3H$.

Claim 6 (previously presented): A compound according to claim 1 wherein Z^1 and Z^2 are selected from phenyl and naphthyl moieties.

Claims 7-9 (cancelled)


Claim 10 (previously presented): A compound according to claim 1 wherein said target bonding group F comprises an affinity tag.

Claim 11 (previously presented): A compound according to claim 1 wherein said spacer group E is selected from:



where Q is selected from: $-CHR'-$, $-NR'-$, $-O-$, $-CR'=CR'-$, $-C(O)-NR'-$ and $-C(O)-O-$; R' is hydrogen or C₁ – C₄ alkyl, p is 0 – 5 and r is 1 – 5.

Claim 12 (original): A compound according to claim 11 wherein Q is selected from:

–CHR', –C(O)–NH– and ; where R' is hereinbefore defined.


Claim 13 (previously presented): A compound according to claim 1 wherein said group –E– F comprises a carboxypentyl group.

Claim 14 (previously presented): A compound according to claim 1 selected from:

- i) 2-{(1E,3E,5E)-5-[1-(5-carboxypentyl)-3-methyl-5-sulpho-3-(4-sulphobutyl)-1,3-dihydro-2H-indol-2-ylidene]penta-1,3-dienyl}-1-ethyl-3-methyl-3-(4-sulphobutyl)-3H-indolium-5-sulphonate;
- ii) 2-{(1E,3E,5E)-5-[1-(5-carboxypentyl)-3-methyl-5-sulpho-3-(4-sulphobutyl)-1,3-dihydro-2H-indol-2-ylidene]penta-1,3-dienyl}-3-methyl-1,3-bis(4-sulphobutyl)-3H-indolium-5-sulphonate;
- iii) 2-{(1E,3E,5E,7E)-7-[1-(5-carboxypentyl)-3-methyl-5-sulpho-3-(4-sulphobutyl)-1,3-dihydro-2H-indol-2-ylidene]hepta-1,3,5-trienyl}-1-ethyl-3-methyl-3-(4-sulphobutyl)-3H-indolium-5-sulphonate;
- iv) 2-{(1E,3E,5E,7E)-7-[5-(carboxymethyl)-3-methyl-1,3-bis(4-sulphobutyl)-1,3-dihydro-2H-indol-2-ylidene]hepta-1,3,5-trienyl}-1-ethyl-3-methyl-3-(4-sulphobutyl)-3H-indolium-5-sulphonate; and
- v) 1-benzyl-2-{(1E,3E,5E)-5-[1-(5-carboxypentyl)-3-methyl-5-sulpho-3-(4-sulphobutyl)-1,3-dihydro-2H-indol-2-ylidene]penta-1,3-dienyl}-3-methyl-3-(4-sulphobutyl)-3H-indolium-5-sulphonate.


Claim 15 (withdrawn): A method for preparing a compound according to claim 1, the method comprising:

a) reacting a first intermediate compound having the formula (A):



wherein Z^1 , R^1 , R^3 , R^4 , R^{11} and R^{12} are hereinbefore defined;

b) a second intermediate compound which may be the same or different from the first intermediate compound and having the formula (B):




wherein Z^2 , R^2 , R^5 , R^6 , R^{13} and R^{14} are hereinbefore defined; and

c) a third compound (C) suitable for forming a linkage between the first and second compounds;

provided that at least one of the groups R^1 , R^2 , R^3 , R^4 , R^5 and R^6 is the group $-E-F$, where E and F are hereinbefore defined; and provided that one or more of groups R^{11} , R^{12} , R^{13} and R^{14}

are independently selected from the group $-(\text{CH}_2)_k-\text{W}$, where W is selected from sulphonic acid and phosphonic acid groups and k is an integer from 1 to 10.

Claim 16 (withdrawn): A compound of formula:



wherein:


groups R^3 and R^4 are attached to the Z^1 ring structure, wherein Z^1 is hereinbefore defined; at least one of the groups R^1 , R^3 and R^4 is the group $-\text{E}-\text{F}$ where E and F are hereinbefore defined;

at least one of groups R^{11} and R^{12} is the group $-(\text{CH}_2)_k-\text{W}$, where W is selected from sulphonic acid and phosphonic acid groups and k is an integer from 1 to 10.

Claim 17 (withdrawn): A compound according to claim 16 wherein $-(\text{CH}_2)_k-\text{W}$ is selected from $-(\text{CH}_2)_3-\text{SO}_3\text{H}$ and $-(\text{CH}_2)_4-\text{SO}_3\text{H}$.

Claim 18 (withdrawn): A method for labeling a target component, the method comprising:

- i) contacting said component with a compound of formula (I):



(I)

wherein:

groups R^3 and R^4 are attached to the Z^1 ring structure and groups R^5 and R^6 are attached to the Z^2 ring structure, and $n = 1, 2$ or 3 ;

Z^1 and Z^2 independently represent the carbon atoms necessary to complete a one ring, or two-fused ring aromatic system;

at least one of groups R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 is the group $-E-F$ where E is a single bond or a spacer group having a chain from 1–20 linked atoms selected from the group consisting of carbon, nitrogen and oxygen atoms and F is a target bonding group;


one or more of groups R^{11} , R^{12} , R^{13} and R^{14} are independently selected from the group $-(CH_2)_k-W$, where W is sulphonic acid or phosphonic acid and k is an integer from 1 to 10;

when any of groups R^1 and R^2 is not said group $-E-F$, said remaining groups R^1 and R^2 are independently selected from C_1-C_6 alkyl, benzyl either unsubstituted or substituted with sulphonic acid, and the group $-(CH_2)_k-W$, where W and k are hereinbefore defined;

when any of groups R³, R⁴, R⁵ and R⁶ is not said group –E–F, said remaining groups R³, R⁴, R⁵ and R⁶ are independently selected from hydrogen and sulphonic acid;

when any of groups R¹¹, R¹², R¹³ and R¹⁴ is not said group –(CH₂)_k–W, said remaining groups R¹¹, R¹², R¹³ and R¹⁴ are independently C₁–C₆ alkyl;

remaining groups R⁷ are hydrogen or two of R⁷ together with the group,



form a hydrocarbon ring system having 5 or 6 atoms; and

- ii) incubating said fluorescent dye with said component under conditions suitable for binding to and thereby labeling said component.

Claim 19 (withdrawn): A method according to claim 18 wherein said component is selected from the group consisting of antibody, lipid, protein, peptide, carbohydrate, nucleotides which contain or are derivatized to contain one or more of an amino, sulphhydryl, carbonyl, hydroxyl and carboxyl and thiophosphate groups, and oxy or deoxy polynucleic acids which contain or are derivatized to contain one or more of an amino, sulphhydryl, carbonyl, hydroxyl, carboxyl and thiophosphate groups, microbial materials, drugs, hormones, cells, cell membranes and toxins.